

Painful skin lesions and bloody diarrhea

Abimbola Aderinto-Adike, Bashar Hannawi, Anusha Thomas, Cima Nowbakht, Joseph Varon

Introduction

Pyoderma gangrenosum (PG) is a rare ulcerative lesion, commonly associated with an underlying systemic disease. The diagnosis of pyoderma gangrenosum can be challenging, and often requires a good history and exclusion of other ulcerative cutaneous disease. We present an impressive case of pyoderma gangrenosum in a young woman with bloody diarrhea.

Case presentation

A 21-year-old woman with a distant history of bloody diarrhea one year prior to presentation, came to our institution with a month history of painful skin lesions on her legs and hands. She reported that these lesions started while working in Southern Africa. She denied any fevers, or chills. She reported persistent bloody diarrhea that started about two months prior to onset of her skin lesions. She denied any abdominal pain, weight loss, nausea or vomiting. She had no history of cutaneous trauma. On examination, she was afebrile. Abdomen was soft, non-tender with normal active bowel sounds. A skin lesion on her lower extremity is shown in **Figure 1**.

A biopsy of the lesion showed ulcerated skin bullous lesions with inflammatory cell infiltrates. Colonoscopy revealed diffuse ulceration with gland distortion and an increase in inflammatory cells. The patient was started on prednisone, at 1 mg/kg, which resolved her bloody diarrhea. She was also started on mesalamine to follow up as an outpatient.

Discussion

Pyoderma gangrenosum (PG) is a rare ulcerating

skin lesion that can mimic a wound infection. It is an autoimmune disorder that results in ulceration and necrosis of the skin. Up to 70% of cases with PG are associated with an underlying systemic disease including inflammatory bowel disease, rheumatoid arthritis or hematological malignancies. (1) In patients with IBD, the severity of the disease is not associated with PG but likely indicates a particular phenotype of the disease. (2) Diagnosis is typically made by clinical history and presentation of a rapidly progressive painful ulcer. Histopathology is often non-specific for PG. (3)

Surgical management of PG is controversial as debridement can worsen the inflammatory reaction and exacerbate the skin ulcerations. (4) PG is usually treated with topical steroids or topical calcineurin inhibitors. More extensive disease is usually treated with systemic steroids, immunomodulators such as cyclosporine or newer biologics like infliximab.

Acknowledgment

The authors acknowledge the initial skin and colon pathology slides for this patient were done at Aga Khan Hospital, Nairobi, Kenya.

References

1. Momeni A, Satterwhite T, Eggleston JM 3rd. Postsurgical pyoderma gangrenosum after autologous breast reconstruction: Case report and review of the literature. *Ann Plast Surg* 2013 Oct 16. [Epub ahead of print].
2. Farhi D, Cosnes J, Zizi N, Chosidow O, Seksik P, Beaugerie L, et al. Significance of erythema nodosum and pyoderma gangrenosum in inflammatory bowel diseases: a cohort study of 2402 patients. *Medicine (Baltimore)* 2008;87: 281-93.
3. Agrawal S, Singhanian B. Pyoderma gangrenosum. *BMJ Case Rep* 2010;bcr0420102942.
4. Hill DS, O'Neill JK, Toms A, Watts AM. Pyoderma gangrenosum: A report of a rare complication after knee arthroplasty requiring muscle flap cover supplemented by negative pressure therapy and hyperbaric oxygen. *J Plast Reconstr Aesthet Surg* 2011; 64:1528-32.

From Houston Methodist Hospital, Houston, Texas, USA (Abimbola Aderinto-Adike, Bashar Hannawi, Anusha Thomas, Cima Nowbakht, and Joseph Varon).

Address for correspondence:

Abimbola Aderinto, MD
Houston Methodist Hospital,
6550 Fannin Street, SM1001
Houston, TX 77030, USA

Figure 1. Lesion in lower extremity

